

IMSc/2003/04/05
hep-th/0304152

Asymptotic Density of Open p -brane States with Zero-modes included

S. Kalyana Rama

Institute of Mathematical Sciences, C. I. T. Campus,
Taramani, CHENNAI 600 113, India.

email: krama@imsc.res.in

ABSTRACT

We obtain the asymptotic density of open p -brane states with zero-modes included. The resulting logarithmic correction to the p -brane entropy has a coefficient $-\frac{p+2}{2p}$, and is independent of the dimension of the embedding spacetime. Such logarithmic corrections to the entropy, with precisely this coefficient, appear in two other contexts also: a gas of massless particles in p -dimensional space, and a Schwarzschild black hole in $(p+2)$ -dimensional anti de Sitter spacetime.

1. The asymptotic density of states $\rho(N)$ at level N , $N \gg 1$, for p -branes compactified on $(S^1)^p \times \mathbf{R}^{D-p}$ has been calculated within the semiclassical quantisation scheme [1, 2, 3, 4]. (For various applications of this result see the review article in [4] and [2, 5].) The corresponding p -brane entropy, given by $\ln \rho(N)$, then has a logarithmic correction with a particular coefficient X , which depends on the dimension D of the embedding spacetime.

The correct counting of the asymptotic density of states must also include the zero-mode states. They have been included for the open string case ($p = 1$) in [6, 7]. As a consequence, the logarithmic correction coefficient X becomes independent of D and is given by $X = -\frac{3}{2}$. Such logarithmic corrections to the entropy, with precisely this coefficient, $-\frac{3}{2}$, have appeared in other contexts also: in $(1+1)$ -dimensional conformal field theories [6, 7], and in the entropies for $(2+1)$ and $(3+1)$ dimensional black holes calculated using the spin network formalism [8, 9].

In this paper, using the results of [4], we obtain the asymptotic density of states for open p -branes. We then include the zero-modes following the methods of [6, 7]. We find that the logarithmic correction coefficient X becomes independent of the dimension D of the embedding spacetime, and is given by $X = -\frac{p+2}{2p}$.

Logarithmic corrections to entropy also arise for statistical mechanical systems due to statistical fluctuations [10]. Using the results of [10], we find that logarithmic corrections to the entropy, with precisely the same coefficient as that obtained in the open p -brane case, namely $-\frac{p+2}{2p}$, appear in two other contexts also: a gas of massless particles in p -dimensional space, and a Schwarzschild black hole in $(p+2)$ -dimensional anti de Sitter spacetime [10].

This paper is organised as follows. In section **2**, we briefly present the results of [4] and, using them, obtain the asymptotic density of states for open p -branes. In section **3**, we include the zero-modes. In section **4**, we show that such logarithmic corrections with precisely the same coefficient appear in other contexts also. In section **5**, we conclude by mentioning a few issues for further study.

2. The asymptotic density of states $\rho(N)$ at level N , $N \gg 1$, for p -branes compactified on $(S^1)^p \times \mathbf{R}^{D-p}$ can be calculated within the semiclassical quantisation scheme, and is of the form

$$\rho(N) \simeq C N^B e^{A N^\delta} \quad (1)$$

where δ , A , B , and C are constants. δ was obtained in [1, 2] and the correct expressions for the remaining constants A , B , and C in [4]. The corresponding p -brane entropy $S(N)$ is given by

$$S(N) = \ln \rho(N) \simeq S_0 + X \ln S_0 + (const) \quad (2)$$

where the leading term $S_0 = AN^\delta$ and $X = \frac{B}{\delta}$ is the coefficient of the logarithmic correction to the entropy.

We now present briefly the results of [4]. See [4] for details. In the semiclassical quantisation, the total number operator $\mathcal{N}$ can be written in the proper time formalism as

$$\mathcal{N} = \sum_{i=1}^d \sum_{\mathbf{n} \neq \mathbf{0}} \omega_{\mathbf{n}} \mathcal{N}_{\mathbf{n}}^i, \quad \omega_{\mathbf{n}} \equiv \sqrt{\sum_{j=1}^p n_j^2}, \quad (3)$$

where $d = (D - p - 1)$, $\mathbf{n} = (n_1, n_2, \dots, n_p) \in \mathbf{Z}^p$, $\mathbf{0} = (0, 0, \dots, 0)$, and $\mathcal{N}_{\mathbf{n}}^i$ are number operators [11]. For the sake of simplicity, we have set the p -brane tension to unity and taken all the circles in $(S^1)^p$ to be of unit radius.

Let $\rho(N)$ be the number of independent eigenstates of the total number operator $\mathcal{N}$ with eigenvalue N . Its generating function $F(z)$ is given by

$$F(z) = \sum_{N=0}^{\infty} \rho(N) e^{-zN} = \text{Tr} e^{-z\mathcal{N}} = \prod_{\mathbf{n} \neq \mathbf{0}} (1 - e^{-z\omega_{\mathbf{n}}})^{-d}. \quad (4)$$

Inverting the above relation then gives $\rho(N)$ in terms of $F(z)$:

$$\rho(N) = -\frac{1}{2\pi i} \oint dz e^{Nz} F(z) \quad (5)$$

where the integration contour is a small circle around the origin. Using Meinardus theorem [12] and the properties of Epstein zeta function, an asymptotic expression for $F(z)$ can be obtained in the limit $\text{Re}(z) \rightarrow 0$ which is sufficient to obtain $\rho(N)$ in the limit $N \gg 1$. The asymptotic expression for $F(z)$ is of the form [4]

$$F(z) \simeq c z^b e^{az^{-p}} \quad (6)$$

where a , b , and c are constants. The contour integral in (5), and thus $\rho(N)$, can then be evaluated by saddle point method. For $F(z)$ of the form given

in equation (6), the saddle point is located at $z = z_0 = \left(\frac{N}{ap}\right)^{-\frac{1}{p+1}}$ and $\rho(N)$, in the limit $N \gg 1$, is given by

$$\rho(N) \simeq \frac{c (ap)^{\frac{2b+1}{2(p+1)}}}{\sqrt{2\pi(p+1)}} N^{-\frac{2b+p+2}{2(p+1)}} e^{AN^{\frac{p}{p+1}}}, \quad A = \frac{p+1}{p} (ap)^{\frac{1}{p+1}}. \quad (7)$$

In our case, $a = \frac{2d \Gamma(p) \pi^{\frac{p}{2}} \zeta(p+1)}{\Gamma(\frac{p}{2})}$ and $b = d$ where $d = (D - p - 1)$ and ζ is the Riemann zeta function; the constant c is given explicitly in [4] and is not required for our purposes. See [4] for further details.

We now obtain the asymptotic density of states for an open p -brane using the above results. In this case modes $\mathbf{n}$ and $-\mathbf{n}$ together contribute to one standing wave (SW) mode of the open p -brane and, hence, should be counted only once. Thus, the corresponding total number operator is given by

$$\mathcal{N} = \sum_{i=1}^d \sum_{\text{SW}; \mathbf{n} \neq \mathbf{0}} \omega_{\mathbf{n}} \mathcal{N}_{\mathbf{n}}^i. \quad (8)$$

The corresponding generating function $F_o(z)$ is given by

$$F_o(z) = \sum_{N=0}^{\infty} \rho_o(N) e^{-zN} = \prod_{\text{SW}; \mathbf{n} \neq \mathbf{0}} (1 - e^{-z\omega_{\mathbf{n}}})^{-d} = \prod_{\mathbf{n} \neq \mathbf{0}} (1 - e^{-z\omega_{\mathbf{n}}})^{-\frac{d}{2}} \quad (9)$$

where the last equality follows since $\omega_{\mathbf{n}} = \omega_{-\mathbf{n}}$ and the product in the last expression includes both $\mathbf{n}$ and $-\mathbf{n}$. Note that the generating function $F_o(z)$ is identical to that in equation (4) with d there replaced by $\frac{d}{2}$. It therefore follows that the density of states $\rho_o(N)$ for an open p -brane in the limit $N \gg 1$ is given by (7), but now with d replaced by $\frac{d}{2}$. Explicitly, in the limit $N \gg 1$,

$$\rho_o(N) \simeq \frac{c (ap)^{\frac{D-p}{2(p+1)}}}{\sqrt{2\pi(p+1)}} N^{-\frac{D+1}{2(p+1)}} e^{AN^{\frac{p}{p+1}}}, \quad A = \frac{p+1}{p} (ap)^{\frac{1}{p+1}} \quad (10)$$

where $a = \frac{(D-p-1) \Gamma(p) \pi^{\frac{p}{2}} \zeta(p+1)}{\Gamma(\frac{p}{2})}$, c is as given in [4] but with $d = (D - p - 1)$ there replaced by $\frac{d}{2} = \frac{D-p-1}{2}$, and we have used $b = \frac{d}{2} = \frac{D-p-1}{2}$ in obtaining (10). Note that $\rho(N)$ is of the form given in equation (1) with $\delta = \frac{p}{p+1}$ and $B = -\frac{D+1}{2(p+1)}$, and that string theory result [3] is obtained upon setting $p = 1$.

3. We now include the zero-modes and obtain the resulting asymptotic density of open p -brane states. The complete Hamiltonian for a p -brane in the proper time formalism is given by

$$H = \mathbf{p}^2 + \mathcal{N} \quad (11)$$

where $\mathbf{p}^2$ is the transverse momentum square operator and $\mathcal{N}$ is the total number operator given in (8), and they both commute with each other [4].

The correct counting of the total number of states $\rho(N)$ must include the zero-mode states also, namely those corresponding to the transverse momentum. For open strings ($p = 1$), $\rho(N)$ with zero-modes included has been calculated in [6, 7] in the limit $N \gg 1$. The resulting $\rho(N)$ is still given by equation (1) with $\delta = \frac{1}{2}$ as before, but now with $B = -\frac{3}{4}$ independent of the dimension D of the embedding spacetime. See [6, 7] for further discussions.

The total number of states $\rho(N)$ for open p -branes, with zero-modes included, can be calculated for other values of p also in the limit $N \gg 1$. $\rho(N)$ is still given by equation (5), but now the relevant generating function $F(z)$, with zero-modes included, is given by

$$F(z) = \text{Tre}^{-zH} = \left(\int d^{D-p-1} p e^{-z\mathbf{p}^2} \right) \text{Tre}^{-zN} = \left(\int d^{D-p-1} p e^{-z\mathbf{p}^2} \right) F_o(z) \quad (12)$$

where the second equality follows since the operators $\mathbf{p}^2$ and $\mathcal{N}$ commute with each other and $F_o(z)$ is the generating function given in equation (9). The momentum integral can be evaluated easily and results in an extra z -dependent factor given by

$$\int d^{D-p-1} p e^{-z\mathbf{p}^2} = c_0 z^{-\frac{D-p-1}{2}}$$

where c_0 is a constant. The contour integral in (5), and thus $\rho(N)$, can now be evaluated by saddle point method. The saddle point is at $z = z_0 = \left(\frac{N}{ap} \right)^{-\frac{1}{p+1}}$ as before, and the total number of open p -brane states $\rho_o(N)$, in the limit $N \gg 1$ and with zero-modes included, is now given by

$$\rho_o(N) \simeq \frac{cc_0 (ap)^{\frac{1}{2(p+1)}}}{\sqrt{2\pi(p+1)}} N^{-\frac{p+2}{2(p+1)}} e^{AN \frac{p}{p+1}} \quad (13)$$

where the constants a and c are as in equation (10). Clearly, $\rho(N)$ is of the form given in equation (1) with $\delta = \frac{p}{p+1}$ as before, but now with $B = -\frac{p+2}{2(p+1)}$ independent of the dimension D of the embedding spacetime.

The open p -brane entropy $S(N)$ with zero-modes included is thus given by

$$S(N) = \ln \rho(N) \simeq S_0 - \frac{p+2}{2p} \ln S_0 + (\text{const}) \quad (14)$$

where $S_0 = AN^{\frac{p}{p+1}}$. The coefficient of the logarithmic correction to the open p -brane entropy with zero-modes included is now given by $-\frac{p+2}{2p}$ and is independent of the dimension D of the embedding spacetime. Note that without zero-modes included it is given by $-\frac{D+1}{2p}$, as can be seen from equations (2) and (10), and depends on D .

4. For open strings ($p = 1$) with zero-modes included, the coefficient of the logarithmic correction to the entropy given above becomes $-\frac{3}{2}$, which agrees with the results of [6, 7]. Such logarithmic corrections, with precisely this coefficient, $-\frac{3}{2}$, have appeared in other contexts also: in $(1+1)$ -dimensional conformal field theories [6, 7], and in the entropies for $(2+1)$ and $(3+1)$ dimensional black holes calculated using the spin network formalism [8, 9].

Similarly, it turns out that logarithmic corrections, with precisely the coefficient given in equation (14), namely $-\frac{p+2}{2p}$, also appear in two other contexts. Recently, logarithmic corrections to entropies of statistical mechanical systems, arising due to statistical fluctuations, have been obtained in [10]. One calculates the density of states $\rho(E)$ as an inverse Laplace transformation of the partition function in the canonical ensemble. Statistical fluctuations can then be incorporated naturally. Then, $\rho(E)\Delta$ is the number of states with energy in the range $E \pm \frac{\Delta}{2}$ where Δ depends on the precision with which the system is prepared and, in particular, is independent of E . The entropy $S(E)$ is therefore given by $S(E) = \ln \rho(E) + (\text{const})$.

The result of [10] is that for a system at temperature T , with specific heat $\mathcal{C}$ (which must be positive for this formalism to be applicable), one obtains for its entropy

$$S = S_0 - \frac{1}{2} \ln(\mathcal{C} T^2) + (\text{const}) \quad (15)$$

where S_0 is the leading term. See [10] for details.

Now, consider a gas of massless particles in p -dimensional space. Then

$$S_0 \propto T^p, \quad E \propto T^{p+1}, \quad \mathcal{C} \propto T^p.$$

Equation (15), therefore, gives

$$S = S_0 - \frac{p+2}{2p} \ln S_0 + (const). \quad (16)$$

For a Schwarzschild black hole, the specific heat is negative and, hence, the above formalism is inapplicable [10]. However, a Schwarzschild black hole of sufficiently large mass in a d -dimensional anti de Sitter spacetime (AdS_d) has positive specific heat. Consider an AdS_{p+2} Schwarzschild black hole of mass M . For sufficiently large M , one has [13]

$$S_0 \propto r_+^p, \quad E = M \propto r_+^{p+1}, \quad T \propto r_+,$$

where r_+ is the horizon. It then follows that $\mathcal{C} \propto r_+^p$. Equation (15), therefore, gives

$$S = S_0 - \frac{p+2}{2p} \ln S_0 + (const). \quad (17)$$

See [10]) for details. For AdS_3 , see also [8, 9]. From equations (16) and (17), we see that logarithmic corrections to the entropy of a gas of massless particles in p -dimensional space, and to that of an AdS_{p+2} Schwarzschild black hole, both have a coefficient $-\frac{p+2}{2p}$ which is precisely the same as that obtained in the open p -brane case with zero-modes included.

5. To summarise, we have obtained the asymptotic density of open p -brane states with zero-modes included. The corresponding open p -brane entropy has a logarithmic correction, with a coefficient $-\frac{p+2}{2p}$. Such logarithmic corrections, with precisely the same coefficient, also appear for a p -dimensional gas and for an AdS_{p+2} Schwarzschild black hole where the corrections arise due to statistical fluctuations.

The relation of a p -dimensional gas to AdS_{p+2} Schwarzschild black hole, for $p = 1, 2, 3$, and 5, can be understood in the context of AdS/CFT duality [14] as that of a boundary conformal field theory at high temperature [13]. In light of the present results, one may explore the relations between quantum/semi classical p -branes, p -dimensional gas, and AdS_{p+2} spacetime in more detail. In particular, it will be interesting to know if the values of

p are restricted for quantum/semi classical p -branes, or if a duality exists between p -dimensional gas and AdS_{p+2} spacetime for any value of p .

As mentioned earlier, the coefficient $X = -\frac{3}{2}$, corresponding to $p = 1$, also appears for the entropy of a $(3 + 1)$ dimensional black hole calculated using the spin network formalism [8]. In this formalism, one considers punctures, each carrying a spin $\mathbf{J}_{puncture}$, and counts the number of spin singlet states, namely those states with $\mathbf{J}(total) = 0$. The leading term (and a part of the logarithmic correction) in the entropy corresponds to the number of states with $J_z(total) = 0$. However, such states include states with $\mathbf{J}(total) \neq 0$ also. A correct counting, that counts states with $\mathbf{J}(total) = 0$ only, then leads to the coefficient $-\frac{3}{2}$ for the logarithmic correction to the entropy [8]. It will be interesting to find if a similar interpretation exists for the logarithmic correction coefficient $X = -\frac{p+2}{2p}$ for other values of p also.

Acknowledgement: We thank C. Castro for suggesting this problem, pointing out the review article in [4], and for discussions; R. Kaul for discussions regarding reference [7]; and K. Kirsten for correspondence.

References

- [1] S. Fubini, A. J. Hanson, and R. Jackiw, Phys. Rev. **D 7** (1973) 1732.
- [2] A. Strumia and G. Venturi, Lett. Nuovo Cimento **13** (1975) 337; E. Alvarez and T. Ortin, Mod. Phys. Lett. **A 7** (1992) 2889; B. Harms and Y. Leblanc, Phys. Rev. **D 47** (1993) 2438, hep-th/9208070.
- [3] M. B. Green, J. Schwarz, and E. Witten, *Superstring Theory*, section 2.3.5, Cambridge University Press, 1987.
- [4] A. A. Bytsenko, K. Kirsten, and S. Zerbini, Phys. Lett. **B 304** (1993) 235; A. A. Bytsenko, K. Kirsten, and S. Zerbini, Mod. Phys. Lett. **A 9** (1994) 1569, hep-th/9405090. See also the review A. A. Bytsenko, G. Cognola, L. Vanzo, and S. Zerbini, Phys. Rept. **266** (1996) 1, hep-th/9505061 and references therein.
- [5] A. A. Bytsenko and S. D. Odintsov, Prog. Theor. Phys. **98** (1997) 987, hep-th/9611151; A. A. Bytsenko, A. E. Goncalves, and S. D. Odintsov, JETP Lett. **66** (1997) 11, hep-th/9708136.

- [6] S. Carlip, *Class. Quant. Grav.* **17** (2000) 4175, gr-qc/0005017. See also P. Di Francesco, P. Mathieu, and D. Senechal, *Conformal Field Theory*, sections 6.3 and 10.2, Springer 1997.
- [7] R. K. Kaul, hep-th/0302170.
- [8] R. K. Kaul and P. Majumdar, *Phys. Lett. B* **439** (1998) 267, gr-qc/9801080; *Phys. Rev. Lett.* **84** (2000) 5255, gr-qc/0002040; S. Das, R. K. Kaul, and P. Majumdar, *Phys. Rev. D* **63** (2001) 044019, gr-qc/0006211; R. K. Kaul and S. Kalyana Rama, to appear in *Phys. Rev. D*, gr-qc/0301128.
- [9] D. Birmingham and S. Sen, *Phys. Rev. D* **63** (2001) 047501, hep-th/0008051; T. R. Govindarajan, R. K. Kaul, and V. Suneeta, *Class. Quant. Grav.* **18** (2001) 2877, gr-qc/0104010.
- [10] S. Das, P. Majumdar, and R. K. Bhaduri, *Class. Quant. Grav.* **19** (2002) 2355, hep-th/0111001.
- [11] For the definition of the number operators $\mathcal{N}_{\mathbf{n}}^i$ and their commutation properties see, for example, M. J. Duff et al, *Nucl. Phys. B* **297** (1988) 515.
- [12] See, for example, G. E. Andrews, *The Theory of Partitions*, Chapter 6, *Encyclopedia of Mathematics and its Applications*, Volume 2, Addison-Wesley publishing company (1976).
- [13] E. Witten, *Adv. Theor. Math. Phys.* **2** (1998) 505, hep-th/9803131.
- [14] J. M. Maldacena, *Adv. Theor. Math. Phys.* **2** (1998) 231, hep-th/9711200; S. S. Gubser, I. R. Klebanov, and A. M. Polyakov, *Phys. Lett. B* **428** (1998) 105, hep-th/9802109; E. Witten, *Adv. Theor. Math. Phys.* **2** (1998) 253, hep-th/9802150.